

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/8454078>

Sirtuin activators mimic caloric restriction and delay ageing in metazoans

Article in *Nature* · September 2004

DOI: 10.1038/nature02789 · Source: PubMed

CITATIONS

1,506

READS

552

7 authors, including:



Konrad Howitz

Reaction Biology Corp.

32 PUBLICATIONS 7,595 CITATIONS

[SEE PROFILE](#)



David A Sinclair

Harvard University

301 PUBLICATIONS 44,266 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Plant cell cultures for the production of health promoting metabolites [View project](#)



NAMPT overexpression in mouse skeletal muscle to improve metabolic health [View project](#)

24. Culling, J. F. & Summerfield, Q. Measurements of the binaural temporal window using a detection task. *J. Acoust. Soc. Am.* **103**, 3540–3553 (1998).

Acknowledgements We thank C. Rasmussen for the algorithm 'Minimize'. This work was supported by a MRC Career Establishment Grant to D.M. and a MRC Bioinformatics Studentship to N.S.H.

Competing interests statement The authors declare that they have no competing financial interests.

Correspondence and requests for materials should be addressed to D.M. (d.mcalpine@ucl.ac.uk).

Sirtuin activators mimic caloric restriction and delay ageing in metazoans

Jason G. Wood¹*, Blanka Rogina²*, Siva Lavu¹, Konrad Howitz³, Stephen L. Helfand², Marc Tatar⁴ & David Sinclair¹

¹Department of Pathology, Harvard Medical School, 77 Ave. Louis Pasteur, Boston, Massachusetts 02115, USA

²Department of Genetics and Developmental Biology, University of Connecticut Health Center, 263 Farmington Ave., Farmington, Connecticut 06030, USA

³BIOMOL Research Laboratories, Inc., 5120 Butler Pike, Plymouth Meeting, Pennsylvania 19462, USA

⁴Department of Ecology and Evolutionary Biology, Box G-W, Brown University, Providence, Rhode Island 02912, USA

* These authors contributed equally to this work

Caloric restriction extends lifespan in numerous species. In the budding yeast *Saccharomyces cerevisiae* this effect requires Sir2 (ref. 1), a member of the sirtuin family of NAD⁺-dependent deacetylases^{2,3}. Sirtuin activating compounds (STACs) can promote the survival of human cells and extend the replicative lifespan of yeast⁴. Here we show that resveratrol and other STACs activate sirtuins from *Caenorhabditis elegans* and *Drosophila melanogaster*, and extend the lifespan of these animals without reducing fecundity. Lifespan extension is dependent on functional Sir2, and is not observed when nutrients are restricted. Together these data indicate that STACs slow metazoan ageing by mechanisms that may be related to caloric restriction.

Sir2-like proteins (sirtuins) are a family of NAD⁺-dependent deacetylases conserved from *Escherichia coli* to humans^{5–9} (Fig. 1a) that play important roles in gene silencing, DNA repair, rDNA recombination and ageing in model organisms^{2,10–12}. When diet is restricted (caloric restriction), lifespan is extended in diverse species, suggesting that there is a conserved mechanism for nutrient regulation of ageing^{13–17}. In budding yeast, extra copies of SIR2 extend lifespan by 30%, apparently by mimicking caloric restriction^{1,18}. We recently described a group of compounds (STACs) that stimulate the catalytic activity of yeast and human sirtuins, and extend the replicative lifespan of yeast cells by up to 60% (ref. 4).

To establish whether STACs could activate sirtuins from multicellular animals, we developed a cell-based deacetylation assay for *D. melanogaster* S2 cells. Unlike other classes of deacetylases, the sirtuins are insensitive to the inhibitor trichostatin A (TSA). Several classes of polyphenolic STACs, including chalcones, flavones and stilbenes, increased the rate of TSA-insensitive deacetylation (Fig. 1b). To determine whether this activity was due to direct stimulation of a Sir2 homologue, we purified recombinant SIR-2.1 of *C. elegans* and Sir2 of *D. melanogaster* and determined the effect of various STACs on enzymatic activity *in vitro* (Fig. 1c, d). In a

dose-dependent manner, resveratrol stimulated deacetylation up to 2.5-fold for SIR-2.1 (Fig. 1e) and 2.4-fold for Sir2 (Fig. 1f). As previously observed with the yeast and human Sir2 enzymes, resveratrol lowered the K_m of SIR-2.1 for the co-substrate NAD⁺ (Fig. 1g).

Because resveratrol can significantly extend replicative lifespan in yeast⁴, we asked whether STACs could also extend lifespan in the metazoans *C. elegans* and *D. melanogaster*. Wild-type worms were transferred to plates containing 0 or 100 μ M of resveratrol shortly after reaching adulthood. Lifespan was extended up to 14%, using




Figure 1 Effect of polyphenolic STACs on sirtuins. **a**, Sir2 polypeptides from various species. The NAD⁺-binding pocket (grey), substrate-binding groove (black) and per cent homology to SIRT1 are shown. **b**, Effect of polyphenolic STACs (500 μ M) on TSA-insensitive deacetylase activity in *Drosophila* S2 cells. **c**, Fold stimulation of SIR-2.1 by STACs (10 μ M). **d**, Fold stimulation of *Drosophila* Sir2 by STACs (10 μ M). Values are the mean of at least three determinations (\pm s.e.). **e**, **f**, Activation of *C. elegans* SIR-2.1 (**e**) and *Drosophila* Sir2 (**f**) by resveratrol (\pm s.e.). **g**, SIR-2.1 initial rate as a function of NAD⁺ concentration (\pm s.e.). AFU, arbitrary fluorescence units.




Figure 2 *C. elegans* survival on resveratrol. **a**, Survivorship of adult wild-type N2 *C. elegans* treated with resveratrol fed with heat-killed OP50 *E. coli*. **b**, Survivorship of *sir-2.1* mutants treated with resveratrol fed with heat-killed OP50. Adult lifespan of *sir-2.1* animals does not differ significantly from N2 controls. **c**, Survivorship of wild-type N2 *C. elegans* on 100 μ M resveratrol fed with live OP50. **d**, Survivorship of *sir-2.1* mutants on 100 μ M resveratrol fed with live OP50. **e**, Fecundity of adult hermaphrodites on 100 μ M resveratrol (\pm s.d.). **f**, Feeding rates of L4 larval and adult hermaphrodites treated with 100 μ M resveratrol (\pm s.d.).

either heat-killed or live *E. coli* as food supply (Fig. 2a, c, respectively; Supplementary Table ST1; averaged across trials at 100 μ M resveratrol, lifespan was extended 10% on live and dead *E. coli*), and mortality was decreased across all adult ages (Supplementary Fig. S1). To test whether the lifespan extension depends on functional SIR-2.1, we constructed a *sir-2.1* null mutant. The lifespan of this strain was not appreciably shorter than the wild-type N2 control, and adults treated with resveratrol did not exhibit a significant lifespan extension relative to untreated worms (Fig. 2b, d; Supplementary Table ST1). There was no decrease in fecundity associated with resveratrol treatment (Fig. 2e). To rule out the possibility that resveratrol was causing the animals to eat less, thereby inducing a caloric restriction effect indirectly, we measured feeding rates of both L4 larval and adult worms with or without resveratrol and found no differences (Fig. 2f). Throughout the lifespan trials, we used the reproductive suppressant FUDR to prevent accumulation of progeny on the treatment plates.

We also tested whether STACs could extend lifespan in *D. melanogaster* using the standard laboratory wild-type strain Canton-S and normal fly culturing conditions (vials), and a *yw* marked wild-type strain and demographic culturing conditions (cages) (Supplementary Table ST2). Across independent tests in males and females on abundant diet, lifespan was extended up to 23% with fisetin and up to 29% with resveratrol (Fig. 3a, c, e; Supplementary Table ST2; averaged across these trials with 100 μ M resveratrol, lifespan was extended 20% in females and 16% in males). Increased longevity was associated with reduced mortality before day 40 (Supplementary Fig. S1). A restricted diet increased lifespan by 40% in females and by 14% in males (averaged across trials), and under these conditions neither resveratrol nor fisetin further increased longevity (Fig. 3b, d, f). The lack of an additive effect of resveratrol and a low calorie diet suggests that resveratrol extends lifespan through a mechanism related to caloric restriction.




Figure 3 Survival of wild-type female *D. melanogaster* adults fed resveratrol or fisetin. **a**, Canton-S on 15% SY media. **b**, Canton-S on 5% SY media. **c**, *yw* on 3% CSY media. **d**, *yw* on 2% CSY media. **e**, *yw* on 3% CSY media. **f**, *yw* on 2% CSY media. **g**, Mean 5 day

fecundity per female of Canton-S on 15% SY media \pm resveratrol. **h**, Proportion of *yw* females feeding on diet \pm resveratrol in crop-filling assay. **i**, Mean body mass of Canton-S flies on diet \pm resveratrol (10 μ M). Errors represent s.e.




Figure 4 Survival of *D. melanogaster* adults with mutant alleles of *Sir2* fed resveratrol (100 μ M). Females (**a**) and males (**b**) with loss-of-function genotype *Sir2*^{4.5}/*Sir2*^{5.26}. Females (**c**) and males (**d**) with strong hypomorphic genotype *Sir2*¹⁷/*Sir2*^{KG00871}.

Surprisingly, while diet manipulations that extend *D. melanogaster* longevity typically reduce fecundity^{19,20}, longevity-extending doses of resveratrol modestly increased egg production (10 μ M resveratrol: 69.8 eggs/5 days, s.e. = 2.2; control: 59.9 eggs/5 days, s.e. = 2.2; $t = 3.17$, $P = 0.0017$), particularly in the earliest days of adult life (Fig. 3g). The increase in egg production suggests that the lifespan extending effect of resveratrol in *D. melanogaster* was not due to caloric restriction induced by food aversion or lack of appetite. Consistent with this, no decrease in food uptake was seen with resveratrol-fed flies (Fig. 3h). Furthermore, resveratrol-fed flies maintained normal weight (Fig. 3i), except during days 3–6 when resveratrol fed females were laying significantly more eggs than control fed females.

To determine whether resveratrol extends fly lifespan in a *Sir2*-dependent manner, we analysed a *Sir2* allelic series with increasing amounts of *Sir2*. Adult offspring from crosses between independently derived alleles of *Sir2* were tested. Resveratrol failed to extend lifespan in flies completely lacking functional *Sir2* (*Sir2*^{4.5}/*Sir2*^{5.26}) (Fig. 4a, b) or in flies in which *Sir2* is severely decreased (*Sir2*¹⁷/*Sir2*^{KG00871}) (Fig. 4c, d). Resveratrol increased longevity a small but statistically significant amount in flies homozygous for a hypomorphic *Sir2* allele (*Sir2*^{KG0087}/*Sir2*^{KG0087}) (Supplementary Table ST2, trial 6) and increased lifespan up to 17% in flies with one copy of the hypomorphic allele and one copy of a wild-type *Sir2* (Canton-S/*Sir2*^{KG0087}) (Supplementary Table ST2, trial 7). These data suggest that the ability of resveratrol to extend fly lifespan requires functional *Sir2*.

We previously reported that STACs extend the replicative lifespan of yeast cells by mimicking caloric restriction⁴. Here we show that STACs can extend lifespan in *C. elegans* and *D. melanogaster*, both of which are composed of primarily non-dividing (post-mitotic) cells as adults, and whose somatic and reproductive ageing are indepen-

dent measures of senescence. In both species, resveratrol increases lifespan in a *Sir2*-dependent manner and, at least for the fly, this action appears to function through a pathway related to caloric restriction.

Our observation that resveratrol can increase longevity without an apparent cost of reproduction is counter to prevalent concepts of senescence evolution. However, STACs may still entail trade-offs that involve other traits, or that occur only under some environmental conditions^{21,22}. Plants synthesize STACs such as resveratrol in response to stress and nutrient limitation²³, possibly to activate their own sirtuin pathways⁴. These molecules may activate animal sirtuins because they serve as plant defence mechanisms against consumers or because they are ancestrally orthologous to endogenous activators within metazoans. Alternatively, animals may use plant stress molecules as a cue to prepare for a decline in their environment or food supply⁴. Understanding the adaptive significance, endogenous function and evolutionary origin of sirtuin activators will lead to further insights into the underlying mechanisms of longevity regulation, and could aid in the development of interventions that provide the health benefits of caloric restriction. □

Methods

C. elegans media, strains, lifespan and feeding assays

Bristol N2 (*Caenorhabditis Genetics Center*) was used as the wild-type strain. The *sir-2.1* mutant strain was generated by backcrossing VC199 (*sir-2.1(ok434)*) to N2 four times. For the lifespan assays, synchronized animals were transferred to treatment plates as young adults (2 days after hatching, day 0 of assay), and were transferred to fresh treatment plates every 2 days for the first 6 to 8 days of the assay. Treatment plates were standard NGM media with the reproductive suppressant FUDR (Sigma; 100 μ g l⁻¹) containing resveratrol or solvent (DMSO, which does not affect lifespan) added either directly into the agar before pouring (for live OP50 trials) or diluted into PBS and added to the surface of a dry plate to the indicated final concentration (for dead OP50 trials). For heat-killed *E. coli* trials, OP50 cultures were heated to 65 °C for 30 min, then pelleted and resuspended in

1/10 volume in S Basal supplemented with 10 mM MgSO₄. For *E. coli* killed by ampicillin and kanamycin, overnight cultures of OP50 were treated with 500 µg ml⁻¹ ampicillin and 250 µg ml⁻¹ kanamycin overnight at 37 °C. In all assays, worms were monitored daily for mortality by gently probing with a platinum pick. Assays were performed at 24 °C.

To assay worm feeding rates, worms at the indicated stages were placed on treatment plates (no FUDR) for 4–5 h, then videoed for 1 min using a Pixelink PL-662 camera. The frame rate was slowed and the pumping rate of the pharynx was counted. To assay fecundity, gravid hermaphrodites (5 per plate, raised from synchronized L1s on normal or treatment plates) were allowed to lay eggs on their respective media for 5 h, and the total number of eggs was counted.

D. melanogaster media, strains, feeding assay and lifespan assays

Survival assays were conducted independently with adult *D. melanogaster* in two laboratories. At Brown University, all trials used an *yw* marked wild-type strain. Larvae were reared on standard cornmeal-sugar-yeast (CSY) agar diet (cornmeal 5%, sucrose 10.5%, SAF yeast 2% and agar 0.7%). Newly eclosed adults were placed in 1-litre demography cages with approximately 75 males and 75 females. Three to four replicate 1-litre demography cages were used for each treatment group in each trial. Every two days, dead flies were removed and scored, and food vials were replenished. Food vials contained CSY diet with SAF yeast as either 2% or 3% by weight. Test compounds in 100 µl of EtOH (or blank EtOH in controls) were mixed into melted aliquots of the adult food media to make a final concentration of 0, 10 or 100 µM. Fresh stock solutions and adult media were prepared weekly. At the University of Connecticut Health Center, lifespan trials were conducted with the wild type strain Canton-S, *Sir2*^{4,5} and *Sir2*^{5,26} (S. Smolik), *Sir2*²⁷ (S. Astrom) and *Sir2*^{KG00871} (Drosophila Stock Center). Larvae for all tests were reared on standard CSY diet. Newly eclosed adults were incubated in plastic shell vials containing 5 ml of 15% sugar-yeast diet (15% SY) or 5% sugar-yeast (5% SY) diet (15% SY: 15% yeast, 15% sucrose, 2% agar; 5% SY: 5% yeast, 5% sucrose, 2% agar as per ref. 20.). In all trials, ~20 males with ~20 females were placed into each of 10 vials per treatment group. Every two days, flies were passed into new vials and dead flies were counted. Resveratrol in EtOH (or EtOH alone in controls) was added to the media during its preparation after it had cooled to 65 °C and mixed vigorously. Final compound concentrations were 0, 10, 100 or 200 µM. Fresh stock solution and adult media was prepared weekly.

Feeding rate was measured in *yw* females with the crop-filling assay²⁴. Females were held overnight with water and placed on 2% CSY diet containing food colour (FDA Blue 1) and 0, 10 or 100 µM resveratrol with EtOH. The presence of dye-marked food in the crop was scored in sets of 20 females across five 5-min intervals. For body mass measurements, 10 vials with 20 males and 20 females each of wild type CS-5 flies were kept on 15% SY diet with EtOH or with resveratrol in EtOH (10 µM). Males and females were weighed daily.

Sirtuin purification and deacetylation assays

These are described in Supplementary Methods.

Received 13 April; accepted 28 June 2004; doi:10.1038/nature02789.

Published online 14 July 2004.

- Lin, S. J., Defossez, P. A. & Guarente, L. Requirement of NAD and SIR2 for life-span extension by calorie restriction in *Saccharomyces cerevisiae*. *Science* **289**, 2126–2128 (2000).
- Gasser, S. C. M. The molecular biology of the SIR proteins. *Gene* **279**, 1–16 (2001).
- Hekimi, S. & Guarente, L. Genetics and the specificity of the aging process. *Science* **299**, 1351–1354 (2003).
- Howitz, K. T. et al. Small molecule activators of sirtuins extend *Saccharomyces cerevisiae* lifespan. *Nature* **425**, 191–196 (2003).
- Landry, J. et al. The silencing protein SIR2 and its homologs are NAD-dependent protein deacetylases. *Proc. Natl Acad. Sci. USA* **97**, 5807–5811 (2000).
- Imai, S., Armstrong, C. M., Kaerberlein, M. & Guarente, L. Transcriptional silencing and longevity protein Sir2 is an NAD-dependent histone deacetylase. *Nature* **403**, 795–800 (2000).
- Smith, J. S. et al. A phylogenetically conserved NAD⁺-dependent protein deacetylase activity in the SIR2 protein family. *Proc. Natl Acad. Sci. USA* **97**, 6658–6663 (2000).
- Tanner, K. G., Landry, J., Sternglanz, R. & Denu, J. M. Silent information regulator 2 family of NAD-dependent histone/protein deacetylases generates a unique product, 1-O-acetyl-ADP-ribose. *Proc. Natl Acad. Sci. USA* **97**, 14178–14182 (2000).
- Tanny, J. C., Dowd, G. J., Huang, J., Hilz, H. & Moazed, D. An enzymatic activity in the yeast Sir2 protein that is essential for gene silencing. *Cell* **99**, 735–745 (1999).
- Guarente, L. Sir2 links chromatin silencing, metabolism, and aging. *Genes Dev.* **14**, 1021–1026 (2000).
- Tissenbaum, H. A. & Guarente, L. Increased dosage of a sir-2 gene extends lifespan in *Caenorhabditis elegans*. *Nature* **410**, 227–230 (2001).
- Rogina, B., Helfand, S. L. & Frankel, S. Longevity regulation by *Drosophila* Rpd3 deacetylase and caloric restriction. *Science* **298**, 1745 (2002).
- Jiang, J. C., Jaruga, E., Repnevskaya, M. V. & Jazwinski, S. M. An intervention resembling caloric restriction prolongs life span and retards aging in yeast. *FASEB J.* **14**, 2135–2137 (2000).
- Kenyon, C. A conserved regulatory mechanism for aging. *Cell* **105**, 165–168 (2001).
- Masoro, E. J. Caloric restriction and aging: an update. *Exp. Gerontol.* **35**, 299–305 (2000).
- Koubova, J. & Guarente, L. How does calorie restriction work? *Genes Dev.* **17**, 313–321 (2003).
- Sinclair, D. A. Paradigms and pitfalls of yeast longevity research. *Mech. Ageing Dev.* **123**, 857–867 (2002).
- Kaerberlein, M., McVey, M. & Guarente, L. The SIR2/3/4 complex and SIR2 alone promote longevity in *Saccharomyces cerevisiae* by two different mechanisms. *Genes Dev.* **13**, 2570–2580 (1999).
- Chippindale, A. K., Leroy, A. M., Kim, S. B. & Rose, M. R. Phenotypic plasticity and selection in *Drosophila* life-history evolution. *J. Evol. Biol.* **6**, 171–193 (1993).
- Chapman, T. & Partridge, L. Female fitness in *Drosophila melanogaster*: an interaction between the effect of nutrition and of encounter rate with males. *Proc. R. Soc. Lond. B* **263**, 755–759 (1996).
- Walker, D. W., McColl, G., Jenkins, N. L., Harris, J. & Lithgow, G. J. Evolution of lifespan in *C. elegans*. *Nature* **405**, 296–297 (2000).
- Marden, J. H., Rogina, B., Montooth, K. L. & Helfand, S. L. Conditional tradeoffs between aging and

organismal performance of Indy long-lived mutant flies. *Proc. Natl Acad. Sci. USA* **100**, 3369–3373 (2003).

23. Soles, G. J., Diamandis, E. P. & Goldberg, D. M. Resveratrol: a molecule whose time has come? And gone? *Clin. Biochem.* **30**, 91–113 (1997).
24. Edgecomb, R. S., Harth, C. E. & Schneiderman, A. M. Regulation of feeding behavior in adult *Drosophila melanogaster* varies with feeding regime and nutritional state. *J. Exp. Biol.* **197**, 215–235 (1994).

Supplementary Information accompanies the paper on www.nature.com/nature.

Acknowledgements We thank S. Parkhurst for the dSir2-pRSETc plasmid, and J. Whetstone, H. Yang and Y. Shi for advice and reagents. *C. elegans* strain VC199 (*sir-2.1(ok434)*) was generated by the *C. elegans* Reverse Genetics Core Facility at the University of British Columbia. This work was supported by the National Institute on Aging, the Harvard-Armstrong Foundation, the Donaghy Foundation, and a generous gift from Harmon Rasnow. S.L.H. is an Ellison Medical Research Foundation Senior Investigator, and D.S. and M.T. are Ellison Medical Research Foundation Fellows. J.W. is supported by an NSF Graduate Research Fellowship.

Competing interests statement The authors declare competing financial interests: details accompany the paper on www.nature.com/nature.

Correspondence and requests for materials should be addressed to D.S. (david_sinclair@hms.harvard.edu), S.L.H. (shelfand@neuron.uchc.edu) or M.T. (marc_tatar@brown.edu).

Planar cell polarity signalling controls cell division orientation during zebrafish gastrulation

Ying Gong^{1,2}, Chunhui Mo² & Scott E. Fraser^{1,2}

¹Biological Imaging Center, Beckman Institute, ²Biochemistry and Molecular Biophysics, Division of Biology, California Institute of Technology, Pasadena, California 91125, USA

Oriented cell division is an integral part of pattern development in processes ranging from asymmetric segregation of cell-fate determinants to the shaping of tissues^{1,2}. Despite proposals that it has an important function in tissue elongation^{3,4}, the mechanisms regulating division orientation have been little studied outside of the invertebrates *Caenorhabditis elegans* and *Drosophila melanogaster*¹. Here, we have analysed mitotic divisions during zebrafish gastrulation using *in vivo* confocal imaging and found that cells in dorsal tissues preferentially divide along the animal–vegetal axis of the embryo. Establishment of this animal–vegetal polarity requires the Wnt pathway components Silberblick/Wnt11, Dishevelled and Strabismus. Our findings demonstrate an important role for non-canonical Wnt signalling in oriented cell division during zebrafish gastrulation, and indicate that oriented cell division is a driving force for axis elongation. Furthermore, we propose that non-canonical Wnt signalling has a conserved role in vertebrate axis elongation, orienting both cell intercalation and mitotic division.

Gastrulation in zebrafish starts as the blastoderm begins to cover the yolk cell. Mesendoderm precursors internalize near the blastoderm margin to form an inner blastoderm stratum termed the hypoblast. Cells remaining in the outer stratum constitute the epiblast, which give rise to neural ectoderm on the dorsal side and epidermis on the ventral side^{5,6}. The dorsal epiblast consists of two–three layers of cells, most of which divide twice during gastrulation: once near the start of gastrulation and another in mid gastrulation^{7,8}. Mitotic divisions at different stages appear to be random, oriented along the animal–vegetal axis, or oriented medio-laterally^{7,8}. Neither the full extent nor the mechanisms that regulate division orientation have been explored. Thus, we have characterized the patterns of cell division throughout the depth of the dorsal

6. Gardel, M. L. *et al.* Elastic behavior of cross-linked and bundled actin networks. *Science* **304**, 1301–1305 (2004).
7. Sherman, M. B. *et al.* The three-dimensional structure of the *Limulus* acrosomal process: a dynamic actin bundle. *J. Mol. Biol.* **294**, 139–149 (1999).
8. Schmid, M. F. Cross-correlation and merging of crystallographic reflections derived from cryoelectron micrographs of 3D crystals: application to the *Limulus* acrosomal bundle. *J. Struct. Biol.* **144**, 195–208 (2003).
9. Schmid, M. F., Agris, J. M., Jakana, J., Matsudaira, P. & Chiu, W. Three-dimensional structure of a single filament in the *Limulus* acrosomal bundle: scruin binds to homologous helix–loop–beta motifs in actin. *J. Cell Biol.* **124**, 341–350 (1994).
10. Galkin, V. E. *et al.* The location of ubiquitin in *Lethocerus* arthrin. *J. Mol. Biol.* **325**, 623–628 (2003).
11. Galkin, V. E. *et al.* The bacterial protein SipA polymerizes G-actin and mimics muscle nebulin. *Nature Struct. Biol.* **9**, 518–521 (2002).
12. Galkin, V. E. *et al.* The utrophin actin-binding domain binds F-actin in two different modes: implications for the spectrin superfamily of proteins. *J. Cell Biol.* **157**, 243–251 (2002).
13. Orlova, A. *et al.* Probing the structure of F-actin: cross-links constrain atomic models and modify actin dynamics. *J. Mol. Biol.* **312**, 95–106 (2001).
14. Galkin, V. E., Orlova, A., Lukyanova, N., Wriggers, W. & Egelman, E. H. Actin depolymerizing factor stabilizes an existing state of F-actin and can change the tilt of F-actin subunits. *J. Cell Biol.* **153**, 75–86 (2001).
15. McGough, A., Pope, B., Chiu, W. & Weeds, A. Cofilin changes the twist of F-actin: implications for actin filament dynamics and cellular function. *J. Cell Biol.* **138**, 771–781 (1997).
16. Jiang, W., Baker, M. L., Ludtke, S. J. & Chiu, W. Bridging the information gap: computational tools for intermediate resolution structure interpretation. *J. Mol. Biol.* **308**, 1033–1044 (2001).
17. Kabsch, W., Mannherz, H. G., Suck, D., Pai, E. F. & Holmes, K. C. Atomic structure of the actin:DNase I complex. *Nature* **347**, 37–44 (1990).
18. Holmes, K. C., Popp, D., Gebhard, W. & Kabsch, W. Atomic model of the actin filament. *Nature* **347**, 44–49 (1990).
19. Dominguez, R. & Graceffa, P. Solution properties of TMR-actin: when biochemical and crystal data agree. *Biophys. J.* **85**, 2073–2074 (2003).
20. Graceffa, P. & Dominguez, R. Crystal structure of monomeric actin in the ATP state. Structural basis of nucleotide-dependent actin dynamics. *J. Biol. Chem.* **278**, 34172–34180 (2003).
21. Sablin, E. P. *et al.* How does ATP hydrolysis control actin's associations? *Proc. Natl Acad. Sci. USA* **99**, 10945–10947 (2002).
22. Holmes, K. C., Angert, I., Kull, F. J., Jahn, W. & Schroder, R. R. Electron cryo-microscopy shows how strong binding of myosin to actin releases nucleotide. *Nature* **425**, 423–427 (2003).
23. Borovikov, Y. S. *et al.* Fluorescence depolarization of actin filaments in reconstructed myofibers: the effect of S1 or pPDM-S1 on movements of distinct areas of actin. *Biophys. J.* **86**, 3020–3029 (2004).
24. Otterbein, L. R., Graceffa, P. & Dominguez, R. The crystal structure of uncomplexed actin in the ADP state. *Science* **293**, 708–711 (2001).
25. Egelman, E. H. Actin allostery again? *Nature Struct. Biol.* **8**, 735–736 (2001).
26. Egelman, E. H., Francis, N. & DeRosier, D. J. F-actin is a helix with a random variable twist. *Nature* **298**, 131–135 (1982).
27. Way, M., Sanders, M., Garcia, C., Sakai, J. & Matsudaira, P. Sequence and domain organization of scruin, an actin-cross-linking protein in the acrosomal process of *Limulus* sperm. *J. Cell Biol.* **128**, 51–60 (1995).
28. Bullitt, E. S., DeRosier, D. J., Coluccio, L. M. & Tilney, L. G. Three-dimensional reconstruction of an actin bundle. *J. Cell Biol.* **107**, 597–611 (1988).
29. Sanders, M., Way, M., Sakai, J. & Matsudaira, P. Characterization of the actin crosslinking properties of the scruin-calmodulin complex from the acrosomal process of *Limulus* sperm. *J. Biol. Chem.* **271**, 2651–2657 (1996).
30. Mahadevan, L. & Matsudaira, P. Motility powered by supramolecular springs and ratchets. *Science* **288**, 95–100 (2000).

Supplementary Information accompanies the paper on www.nature.com/nature.

Acknowledgements This research is supported by the NCCR and NIGMS of NIH. We thank M. Baker for assistance in the helixhunter and foldhunter searches, and M. Dougherty for advice on graphical display.

Competing interests statement The authors declare that they have no competing financial interests.

Correspondence and requests for materials should be addressed to W.C. (wah@bcm.tmc.edu). The accession number is EMD-1088.

addendum

Pathways towards and away from Alzheimer's disease

Mark P. Mattson

Nature **430**, 631–639 (2004).

In the list of treatments given in Fig. 3 of this Review Article, I would like to add “Glutamate-receptor modulating agents (**2, 3, 4, 6**)”, where the numbers in bold indicate the sites of action in the pathogenesis pathway. □

corrigendum

Sirtuin activators mimic caloric restriction and delay ageing in metazoans

Jason G. Wood, Blanka Rogina, Siva Lavu, Konrad Howitz, Stephen L. Helfand, Marc Tatar & David Sinclair

Nature **430**, 686–689 (2004).

There are errors in Fig. 4 of this Letter: panels a and d are correct; however, panel c was incorrectly published as a duplicate of panel a, and panel b should have been labelled as panel c instead. The new panel b is shown here. □

